

Differential-difference system related to toroidal Lie algebra

Saburo Kakei *

Department of Mathematical Sciences,
School of Science and Engineering, Waseda University
Ohkubo 3-4-1, Shinjuku-ku, Tokyo 169-8555, Japan

Yasuhiro Ohta

Department of Applied Mathematics,
Faculty of Engineering, Hiroshima University
Kagamiyama 1-4-1, Higashi-hiroshima 739-8527, Japan
E-mail: ohta@kurims.kyoto-u.ac.jp

Abstract

We present a novel differential-difference system in (2+1)-dimensional space-time (one discrete, two continuum), arisen from the Bogoyavlensky's (2+1)-dimensional KdV hierarchy. Our method is based on the bilinear identity of the hierarchy, which is related to the vertex operator representation of the toroidal Lie algebra $\mathfrak{sl}_2^{\text{tor}}$.

1 Introduction and main results

Multi-dimensional generalization of classical soliton equations has been one of the most exciting topic in the field of integrable systems. Among other things, Calogero [1] proposed an interesting example that is a (2+1)-dimensional extension of the Korteweg-de Vries equation,

$$u_t = \frac{1}{4}u_{xxy} + uu_y + \frac{1}{2}u_x \int^x u_y dx. \quad (1)$$

Yu et al. [2] obtained multi-soliton solutions of the (2+1)-dimensional KdV equation (1) by using the Hirota's bilinear method. Let us consider the following Hirota-type equations,

$$(D_x^4 - 4D_x D_{t'})\tau \cdot \tau = 0, \quad (2)$$

$$(D_y D_x^3 + 2D_y D_{t'} - 6D_t D_x)\tau \cdot \tau = 0, \quad (3)$$

*Present address: Department of Mathematics, Rikkyo University, Nishi-ikebukuro 3-34-1, Toshima-ku, Tokyo 171-8501, Japan. E-mail: kakei@rkmath.rikkyo.ac.jp

where we have used the D -operators of Hirota defined as

$$\begin{aligned} D_x D_y \dots f(x, y, \dots) \cdot g(x, y, \dots) \\ \stackrel{\text{def}}{=} \partial_s \partial_t f(x + s, y + t, \dots) g(x - s, y - t, \dots) |_{s, t, \dots = 0}. \end{aligned}$$

We remark that we have introduced auxiliary variables t' that is a hidden parameter in (1). If we set $u = 2(\log \tau)_{xx}$ and use (2) to eliminate $\partial_{t'}$, then one can show that $u = u(x, y, t)$ solves (1).

Bogoyavlensky [3] showed that there is a hierarchy of higher-order integrable equations associated with (1). In the paper [4], Ikeda and Takasaki generalized the Bogoyavlensky's hierarchy from the viewpoint of the Sato's theory of the KP hierarchy [5, 6, 7, 8], and discussed the relationship to toroidal Lie algebras. We note that the relation between integrable hierarchy and toroidal algebras has been discussed also by Billig [9], Iohara, Saito and Wakimoto [10] by using vertex operator representations.

In the present paper, we propose the following differential-difference system that has the same symmetry:

$$\begin{aligned} \partial_t u_k = \Delta_{-k} \left(\frac{\partial_x u_{k+1}}{1 - \exp(-u_{k+1} - u_k)} \right. \\ \left. - \frac{\partial_x u_k}{1 - \exp(u_{k+1} + u_k)} - \frac{1 + \exp(u_{k+1} + u_k)}{1 - \exp(u_{k+1} + u_k)} v_k \right), \end{aligned} \quad (4)$$

$$\begin{aligned} \Delta_{-k} v_k = \frac{\partial_x u_{k+1}}{u_{k+1}} + \frac{\partial_x (u_{k+1} + u_k)}{1 - \exp(u_{k+1} + u_k)} \\ + \frac{\partial_x u_k}{u_k} + \frac{\partial_x (u_k + u_{k-1})}{1 - \exp(u_k + u_{k-1})}, \end{aligned} \quad (5)$$

where Δ_{-k} denotes the backward difference operator $\Delta_{-k} \stackrel{\text{def}}{=} 1 - \exp(-\partial_k)$ ($\Delta_{-k} u_k = u_k - u_{k-1}$). We also show that this system has soliton-type solutions.

2 Lie algebraic derivation of bilinear identity

Here we briefly review the Lie algebraic derivation of the bilinear identity of the Bogoyavlensky's hierarchy [4], which is a generating function of Hirota-type differential equations. We remark that the Lie algebra considered in [4] is bigger than that of present article. Here we don't include the derivations to $\mathfrak{sl}_2^{\text{tor}}$ since those are not needed for our purpose. Owing to this difference, the proof given below may be simpler than that of [4].

The 2-toroidal Lie algebra $\mathfrak{sl}_2^{\text{tor}}$ [11, 12] is the universal central extension of the double loop algebra $\mathfrak{sl}_2 \otimes \mathbb{C}[s, s^{-1}, t, t^{-1}]$, while the affine Lie algebra $\widehat{\mathfrak{sl}}_2$ is the central extension of $\mathfrak{sl}_2 \otimes \mathbb{C}[t, t^{-1}]$. Let A be the ring of Laurent polynomials of two variables s and t . As a vector space, $\mathfrak{sl}_2^{\text{tor}}$ is isomorphic to $\mathfrak{sl}_2 \otimes \mathbb{C}[s, s^{-1}, t, t^{-1}] \oplus \Omega_A/dA$, where Ω_A denotes the module of Kähler differentials of A defined with the canonical derivation $d : A \rightarrow \Omega_A$. We define the Lie algebra structure of $\mathfrak{sl}_2^{\text{tor}}$ by

$$[x \otimes a, y \otimes b] = [x, y] \otimes ab + (x|y) \overline{(da)b} \quad (x, y \in \mathfrak{sl}_2, a, b \in A), \quad (6)$$

$$[\mathfrak{sl}_2^{\text{tor}}, \Omega_A/dA] = 0, \quad (7)$$

where $(x|y)$ denotes the Killing form and $\overline{\cdot} : \Omega_A \rightarrow \Omega_A/dA$ the canonical projection.

In terms of generating series $X_m(z)$ ($X = E, F, H$, $m \in \mathbb{Z}$), $K_m^s(z)$ and $K_m^t(z)$, defined by

$$\begin{aligned} X_m(z) &\stackrel{\text{def}}{=} \sum_{n \in \mathbb{Z}} X \otimes s^n t^m \cdot z^{-n-1}, \\ K_m^s(z) &\stackrel{\text{def}}{=} \sum_{n \in \mathbb{Z}} \overline{s^n t^m \, \text{d} \log s} \cdot z^{-n}, \\ K_m^t(z) &\stackrel{\text{def}}{=} \sum_{n \in \mathbb{Z}} \overline{s^n t^m \, \text{d} \log t} \cdot z^{-n-1}, \end{aligned}$$

the relation (6) can be expressed as

$$\begin{aligned} X_n(z)Y_n(w) &= \frac{1}{z-w} [X, Y]_{m+n}(w) \\ &+ \frac{1}{(z-w)^2} (X|Y) K_{m+n}^s(w) + \frac{m}{z-w} (X|Y) K_{m+n}^t(w) \\ &+ \text{regular as } z \rightarrow w. \end{aligned} \tag{8}$$

There exists a class of representations of $\mathfrak{sl}_2^{\text{tor}}$, which comes directly from that of $\widehat{\mathfrak{sl}}_2$. We consider the space of polynomials,

$$F_y \stackrel{\text{def}}{=} \mathbb{C}[y_j, j \in \mathbb{Z}] \otimes \mathbb{C}[\exp(\pm y_0)],$$

and define the generating series $\varphi(z)$ and $V_m(y; z)$ by

$$\varphi(z) \stackrel{\text{def}}{=} \sum_{n \in \mathbb{Z}} n y_n z^{n-1}, \quad V_m(y; z) \stackrel{\text{def}}{=} \exp \left[m \sum_{n \in \mathbb{Z}} y_n z^n \right].$$

Proposition 1. (cf. [10, 13]) Let (V, π) be a representation of $\widehat{\mathfrak{sl}}_2$ such that $\overline{\text{d} \log s} \mapsto c \cdot \text{id}_V$ for $c \in \mathbb{C}$. Then we can define the representation π^{tor} of $\mathfrak{sl}_2^{\text{tor}}$ on $V \otimes F_y$ such that

$$\begin{aligned} X_m(z) &\mapsto X^\pi(z) \otimes V_m(z), \\ K_m^s(z) &\mapsto c \cdot \text{id}_V \otimes V_m(z), \\ K_m^t(z) &\mapsto c \cdot \text{id}_V \otimes \varphi(z) V_m(z), \end{aligned}$$

where $X = E, F, H$, $m \in \mathbb{Z}$ and $X^\pi(z) \stackrel{\text{def}}{=} \sum_{n \in \mathbb{Z}} \pi(X \otimes s^n) z^{-n-1}$.

Proof. Using the operator product expansion for $\widehat{\mathfrak{sl}}_2$,

$$\begin{aligned} X(z)Y(w) &= \frac{1}{z-w} [X, Y](w) + \frac{1}{(z-w)^2} (X|Y) K \\ &+ \text{regular as } z \rightarrow w, \end{aligned}$$

and the property $V_m(z)V_n(z) = V_{m+n}(z)$, it is straightforward to show that $X_m(z)$ above satisfies (8). The remaining relations can be checked by direct calculations. $\square$

To see the relationship to soliton theory, we shall consider the representation of $\widehat{\mathfrak{sl}}_2$ on the fermionic Fock space [7, 8]. Let ψ_j, ψ_j^* ($j \in \mathbb{Z}$) be free fermions with the canonical anti-commutation relation. In terms of the generating series defined as

$$\psi(\lambda) = \sum_{n \in \mathbb{Z}} \psi_n \lambda^n, \quad \psi^*(\lambda) = \sum_{n \in \mathbb{Z}} \psi_n^* \lambda^{-n},$$

the canonical anti-commutation relation is written as

$$[\psi(\lambda), \psi^*(\mu)]_+ = \delta(\lambda/\mu), \quad [\psi(\lambda), \psi(\mu)]_+ = [\psi^*(\lambda), \psi^*(\mu)]_+ = 0, \quad (9)$$

where $\delta(\lambda) \stackrel{\text{def}}{=} \sum_{n \in \mathbb{Z}} \lambda^n$ is the formal delta-function.

Consider the fermionic Fock space $\mathcal{F}$ with the vacuum vector $|\text{vac}\rangle$ satisfying

$$\psi_j |\text{vac}\rangle = 0 \quad \text{for } j < 0, \quad \psi_j^* |\text{vac}\rangle = 0 \quad \text{for } j \geq 0,$$

and the dual Fock space $\mathcal{F}^*$ with the dual vacuum vector $\langle \text{vac} |$ satisfying

$$\begin{aligned} \langle \text{vac} | \psi_j &= 0 \quad \text{for } j \geq 0, & \langle \text{vac} | \psi_j^* &= 0 \quad \text{for } j < 0, \\ \langle \text{vac} | \text{vac} \rangle &= 1. \end{aligned}$$

As mentioned in [7, 8], a level-1 representation of $\widehat{\mathfrak{sl}}_2$ is given by the elements,

$$:\psi(\lambda)\psi^*(-\lambda): = \sum_{j,n \in \mathbb{Z}} (-1)^j : \psi_{j+n} \psi_j^* : \lambda^n,$$

where $::$ denotes the fermionic normal ordering, $:\psi_i \psi_j^* : \stackrel{\text{def}}{=} \psi_i \psi_j^* - \langle \text{vac} | \psi_i \psi_j^* | \text{vac} \rangle$. Applying Proposition 1, we can construct a representation of $\mathfrak{sl}_2^{\text{tor}}$ on the space $\mathcal{F}_y \stackrel{\text{def}}{=} \mathcal{F} \otimes F_y$ with the vacuum vector $|\text{vac}\rangle^{\text{tor}} \stackrel{\text{def}}{=} |\text{vac}\rangle \otimes 1$.

We now introduce the following operator acting on $\mathcal{F}_y \otimes \mathcal{F}_{y'}$:

$$\Omega^{\text{tor}} \stackrel{\text{def}}{=} \sum_{m \in \mathbb{Z}} \oint \frac{d\lambda}{2\pi i \lambda} \psi(\lambda) V_m(\lambda; y) \otimes \psi^*(\lambda) V_{-m}(\lambda; y').$$

Using the anti-commutation relation (9) and the relation $V_m(y; \lambda) V_n(y; \lambda) = V_{m+n}(y; \lambda)$, we can obtain the following identity by direct calculations:

$$[\Omega^{\text{tor}}, \psi(p) \psi^*(p) V_n(y; p) \otimes 1 + 1 \otimes \psi(p) \psi^*(p) V_n(y'; p)] = 0,$$

which means the action of $\mathfrak{sl}_2^{\text{tor}}$ on $\mathcal{F}_y \otimes \mathcal{F}_{y'}$ commutes with Ω^{tor} . Then it is straightforward to show that

$$\Omega^{\text{tor}} (g |\text{vac}\rangle^{\text{tor}} \otimes g |\text{vac}\rangle^{\text{tor}}) = 0 \quad (10)$$

for $g = \exp(X)$, $X \in \mathfrak{sl}_2^{\text{tor}}$.

To rewrite (10) into bosonic language, we prepare two lemmas:

Lemma 1 (“Boson-Fermion correspondence”). [7, 8] For any $|\nu\rangle \in \mathcal{F}$, we have the following formulas,

$$\begin{aligned}\langle \text{vac} | \psi_0^* \exp(H(\mathbf{x})) \psi(\lambda) | \nu \rangle &= \exp(\xi(\mathbf{x}, \lambda)) \langle \text{vac} | \exp(H(\mathbf{x} - [\lambda^{-1}])) | \nu \rangle, \\ \langle \text{vac} | \psi_{-1} \exp(H(\mathbf{x})) \psi^*(\lambda) | \nu \rangle &= \lambda \exp(-\xi(\mathbf{x}, \lambda)) \langle \text{vac} | \exp(H(\mathbf{x} + [\lambda^{-1}])) | \nu \rangle,\end{aligned}$$

where we have used the following notation,

$$\begin{aligned}\mathbf{x} &= (x_1, x_3, \dots), & H(\mathbf{x}) &\stackrel{\text{def}}{=} \sum_{n=1}^{\infty} \sum_{j \in \mathbb{Z}} x_n \psi_j \psi_{n+j}^*, \\ \xi(\mathbf{x}, \lambda) &\stackrel{\text{def}}{=} \sum_{n=1}^{\infty} x_n \lambda^n, & [\lambda^{-1}] &\stackrel{\text{def}}{=} (1/\lambda, 1/2\lambda^2, 1/3\lambda^3, \dots).\end{aligned}$$

Lemma 2. [9, 10] Let $P(n) = \sum_{j \geq 0} n^j P_j$, where $P_j \in \text{Diff}(z)$ are differential operators that may not depend on z . If

$$\sum_{n \in \mathbb{Z}} z^n P(n) g(z) = 0$$

for some formal series $g(z) = \sum_j g_j z^j$, then

$$P(\epsilon - z \partial_z) g(z) \big|_{z=1} = 0$$

as a polynomial in ϵ .

Define the τ -function as

$$\tau(\mathbf{x}, \mathbf{y}) \stackrel{\text{def}}{=} {}^{\text{tor}} \langle \text{vac} | \exp(H(\mathbf{x})) g | \text{vac} \rangle^{\text{tor}}.$$

From the relation (10), together with Lemma 1 and Lemma 2, we have the following bilinear identity:

$$\begin{aligned}\oint \frac{d\lambda}{2\pi i} \exp(\xi(\mathbf{x} - \mathbf{x}', \lambda)) \tau(\mathbf{x} - [\lambda^{-1}], y_0 + \eta(\check{\mathbf{b}}, \lambda^2), \check{\mathbf{y}} - \check{\mathbf{b}}) \\ \times \tau(\mathbf{x}' + [\lambda^{-1}], y_0 - \eta(\check{\mathbf{b}}, \lambda^2), \check{\mathbf{y}} + \check{\mathbf{b}}) = 0,\end{aligned} \quad (11)$$

where $\check{\mathbf{y}} \stackrel{\text{def}}{=} (y_2, y_4, \dots)$ and $\eta(\check{\mathbf{b}}, \lambda^2) \stackrel{\text{def}}{=} \sum_{n=1}^{\infty} b_{2n} \lambda^{2n}$.

Expanding (11), we can obtain Hirota-type differential equations including (2), (3) ($x_1 = x$, $x_3 = t'$, $y_0 = y$, $y_2 = t$). In this sense, the bilinear identity (11) is a generating function of Hirota-type differential equations of the Bogoyavlensky's hierarchy. The N -soliton solution of (11) is obtained as follows [4]:

$$\begin{aligned}\tau_N(\mathbf{x}, \mathbf{y}) &= \sum_{l=0}^N \sum_{j_1 < \dots < j_l} c_{j_1 \dots j_l} \prod_{m=1}^l a_{j_m} \exp(\eta_{j_m}(\mathbf{x}, \mathbf{y})), \\ \eta_j(\mathbf{x}, \mathbf{y}) &\stackrel{\text{def}}{=} \sum_{n=1}^{\infty} 2p_j^{2n-1} x_{2n-1} + \sum_{n=1}^{\infty} r_j p_j^{2n} y_{2n}, \\ c_{j_1 \dots j_l} &\stackrel{\text{def}}{=} \prod_{1 \leq m < n \leq l} \frac{(p_{j_m} - p_{j_n})^2}{(p_{j_m} + p_{j_n})^2}.\end{aligned} \quad (12)$$

3 Derivation of the differential-difference system

We now apply the Miwa transformation [8, 14],

$$\begin{aligned}\mathbf{x}' &= l[\alpha] + m[\beta] + n[\gamma], \\ \mathbf{x} &= (l+1)[\alpha] + (m+1)[\beta] + (n+1)[\gamma],\end{aligned}\tag{13}$$

to the bilinear identity (11). Here we have used the notation $l[\alpha] = (l\alpha, l\alpha^2/2, l\alpha^3/3, \dots)$. We first consider the case $\mathbf{b} = (b_2, b_4, \dots) = \mathbf{0}$. In this case, the bilinear identity (11) is reduced to that of the ordinary KP hierarchy. Thus we have the Hirota-Miwa equation (or the discrete KP equation),

$$\begin{aligned}&\alpha(\beta - \gamma) \tau(l+1, m, n; \mathbf{y}) \tau(l, m+1, n+1; \mathbf{y}) \\ &+ \beta(\gamma - \alpha) \tau(l, m+1, n; \mathbf{y}) \tau(l+1, m, n+1; \mathbf{y}) \\ &+ \gamma(\alpha - \beta) \tau(l, m, n+1; \mathbf{y}) \tau(l+1, m+1, n; \mathbf{y}) = 0\end{aligned}\tag{14}$$

where $\tau(l, m, n; \mathbf{y})$ denotes

$$\tau(l, m, n; \mathbf{y}) = \tau(\mathbf{x} = l[\alpha] + m[\beta] + n[\gamma], \mathbf{y}).$$

We then consider the time-evolutions with respect to y_0, y_2 . Collecting the coefficient of b_2 in the bilinear identity (11), we have

$$\oint \frac{d\lambda}{2\pi i} \exp(\xi(\mathbf{x} - \mathbf{x}', \lambda)) (D_{y_2} - \lambda^2 D_{y_0}) \times \tau(\mathbf{x}' + [\lambda^{-1}], \mathbf{y}) \cdot \tau(\mathbf{x} - [\lambda^{-1}], \mathbf{y}) = 0.$$

Applying the Miwa transformation (13), we obtain

$$\begin{aligned}&\alpha^2 \beta \gamma (\beta - \gamma) D_{y_2} \tau(l+1, m, n; \mathbf{y}) \cdot \tau(l, m+1, n+1; \mathbf{y}) \\ &+ \alpha \beta^2 \gamma (\gamma - \alpha) D_{y_2} \tau(l, m+1, n; \mathbf{y}) \cdot \tau(l+1, m, n+1; \mathbf{y}) \\ &+ \alpha \beta \gamma^2 (\alpha - \beta) D_{y_2} \tau(l, m, n+1; \mathbf{y}) \cdot \tau(l+1, m+1, n; \mathbf{y}) \\ &= \beta \gamma (\beta - \gamma) D_{y_0} \tau(l+1, m, n; \mathbf{y}) \cdot \tau(l, m+1, n+1; \mathbf{y}) \\ &+ \gamma \alpha (\gamma - \alpha) D_{y_0} \tau(l, m+1, n; \mathbf{y}) \cdot \tau(l+1, m, n+1; \mathbf{y}) \\ &+ \alpha \beta (\alpha - \beta) D_{y_0} \tau(l, m, n+1; \mathbf{y}) \cdot \tau(l+1, m+1, n; \mathbf{y}) \\ &- (\alpha - \beta)(\beta - \gamma)(\gamma - \alpha) \\ &\quad \times D_{y_0} \tau(l, m, n; \mathbf{y}) \cdot \tau(l+1, m+1, n+1; \mathbf{y})\end{aligned}\tag{15}$$

We further impose the condition $\beta = \gamma$. Then the τ -function $\tau(l, m, n; \mathbf{y})$ depends only on $k \stackrel{\text{def}}{=} m - n$, l and $\mathbf{y}$. In this sense, we rewrite

$$\tau(l, m, n; \mathbf{y}) \rightarrow \tau(l, k; \mathbf{y}).$$

Under this condition, the equations (14) and (15) are reduced to

$$\begin{aligned} & 2\alpha \tau(l+1, k; \mathbf{y}) \tau(l, k; \mathbf{y}) \\ & - (\alpha + \beta) \tau(l, k+1; \mathbf{y}) \tau(l+1, k-1; \mathbf{y}) \\ & - (\alpha - \beta) \tau(l, k-1; \mathbf{y}) \tau(l+1, k+1; \mathbf{y}) = 0, \end{aligned} \quad (16)$$

$$\begin{aligned} & \alpha \beta^2 D_{y_2} (2\alpha \tau(l+1, k; \mathbf{y}) \cdot \tau(l, k; \mathbf{y}) \\ & \quad - (\alpha + \beta) \tau(l, k+1; \mathbf{y}) \cdot \tau(l+1, k-1; \mathbf{y}) \\ & \quad - (\alpha - \beta) \tau(l, k-1; \mathbf{y}) \cdot \tau(l+1, k+1; \mathbf{y})) \\ & = D_{y_0} (2(2\beta^2 - \alpha^2) \tau(l+1, k; \mathbf{y}) \cdot \tau(l, k; \mathbf{y}) \\ & \quad - \alpha(\alpha + \beta) \tau(l, k+1; \mathbf{y}) \cdot \tau(l+1, k-1; \mathbf{y}) \\ & \quad - \alpha(\alpha - \beta) \tau(l, k-1; \mathbf{y}) \cdot \tau(l+1, k+1; \mathbf{y})). \end{aligned} \quad (17)$$

Furthermore, we can construct N -soliton solution by applying (13) to (12):

$$\begin{aligned} \tau_N(l, k; y_0, y_2) &= \sum_{l=0}^N \sum_{j_1 < \dots < j_l} c_{j_1 \dots j_l} \prod_{m=1}^l \phi_{j_m}(l, k; y_0, y_2), \\ \phi_j(l, k; y_0, y_2) &\stackrel{\text{def}}{=} a_j \exp(r_j y_0 + r_j p_j^2 y_2) \left(\frac{1 + p_j \alpha}{1 - p_j \alpha} \right)^l \left(\frac{1 + p_j \beta}{1 - p_j \beta} \right)^k, \end{aligned} \quad (18)$$

where $c_{j_1 \dots j_l}$ is the same as the continuum one (12). We remark that the N -soliton τ -function can be written as Wronskian determinant. Using the determinant expression, we can show that both (16) and (17) are reduced to the Plücker relations.

Introducing the variables as

$$\begin{aligned} \partial_t &\stackrel{\text{def}}{=} \frac{2}{\alpha^2} \partial_{y_0} - 2\partial_{y_2}, & \partial_x &\stackrel{\text{def}}{=} \partial_{y_2} - \frac{1}{\beta^2} \partial_{y_0}, \\ u_k(t, x) &\stackrel{\text{def}}{=} \log \left[\left(\frac{\beta - \alpha}{\beta + \alpha} \right)^{1/2} \frac{\tau(l+1, k+1) \tau(l, k)}{\tau(l, k+1) \tau(l+1, k)} \right], \\ v_k(t, x) &\stackrel{\text{def}}{=} \partial_x \log \frac{\tau(l, k+2)}{\tau(l, k)}, \end{aligned}$$

we have the differential-difference equations (4) and (5), which have the N -soliton solution corresponding to the τ -function (18).

4 Concluding remarks

In this paper, we have introduced the differential-difference system (4) and (5), which is related to toroidal Lie algebra $\mathfrak{sl}_2^{\text{tor}}$. Since the symmetry of the toroidal Lie algebra allows us to introduce extra parameters of wave numbers in the soliton solution (i.e., r_j in (12) and (18)), it might be possible to construct some interesting solutions. In particular, we can obtain a class of traveling-wave solutions that has the shape of the character “V” (Figure 1), which is a special case of the two-soliton solutions. The existence of the V-soliton type solution is one of the features of this class of equations.

We note that there exist solutions of the same shape for the (2+1)-dimensional KdV equation (1), and for a (2+1)-dimensional generalization of the nonlinear Schrödinger

(NLS) equation [15] that also has the symmetry of the toroidal Lie algebra $\mathfrak{sl}_2^{\text{tor}}$ [17]. We also remark that Oikawa et al. [16] discussed the propagation of the V-soliton in a two-layer fluid, which is governed by a equation similar to the (2+1)-dimensional NLS equation.

Acknowledgements

The authors acknowledge Doctor Takeshi Ikeda and Professor Kanehisa Takasaki for discussions. The first author is partially supported by the Grant-in-Aid for Scientific Research (No. 12740115) from the Ministry of Education, Culture, Sports, Science and Technology.

References

- [1] F. Calogero, *Lett. Nouvo Cimento* **14** (1975) 443-447.
- [2] Yu Song-Ju, K. Toda, N. Sasa and T. Fukuyama, *J. Physics* **A31** (1998) 3337-3347.
- [3] O.I. Bogoyavlensky, *Russian Math. Surveys* **45**,4 (1990), 1-86.
- [4] T. Ikeda and K. Takasaki, *preprint* (nlin.SI/0004015), to appear in *Internat. Math. Res. Notices*.
- [5] M. Sato, *RIMS Kokyuroku* **439** (1981) 30-46.
- [6] M. Sato and Y. Sato, in *Nonlinear Partial Differential Equations in Applied Sciences*, (P.D. Lax, H. Fujita, and G. Strang, eds.), 259-271, Kinokuniya/North-Holland, 1982.
- [7] E. Date, M. Jimbo, M. Kashiwara and T. Miwa, in *Proceedings of RIMS symposium Nonlinear integtable systems —classical theory and quantum theory—* (M. Jimbo and T. Miwa, eds.) 39-120, World Scientific, 1983.
- [8] M. Jimbo and T. Miwa, *Publ. RIMS* **19** (1983) 943.
- [9] Y. Billig, *J. Algebra* **217** (1999) 40-64.
- [10] K. Iohara, Y. Saito and M. Wakimoto, *Phys. Lett.* **A254** (1999) 37-46; *ibid.* **A268** (2000) 448-449; *Progr. of Theor. Phys. Suppl.* **135** (1999) 166-181.
- [11] C. Kassel, *J. Pure Appl. Algebra* **34** (1985) 265-275.
- [12] R.V. Moody, S. Eswara Rao and T. Yokonuma, *Geom. Ded.* **35** (1990) 283-307.
- [13] S. Berman and Y. Billig, *J. Algebra* **221** (1999) 188-231.
- [14] T. Miwa, *Proc. Japan Acad.* **58**, Ser. A (1982) 9-12.
- [15] I.A.B. Strachan, *J. Math. Phys.* **34** (1993) 243-259.
- [16] M. Oikawa, M. Okamura and M. Funakoshi, *J. Phys. Soc. Jpn.* **58** (1989) 4416-4430.
- [17] S. Kakei, T. Ikeda and K. Takasaki, in preparation.




Figure 1: Special case of the two-soliton solutions (“V-soliton”) with $p_1 = p_2$ ($p_1 = p_2 = 0.3$, $r_1 = 0.15$, $r_2 = -0.1$, $a_1 = a_2 = 1$, $\alpha = 0.8$, $\beta = 0.5$).




Figure 2: Two-soliton solutions with $p_1 \neq p_2$ ($p_1 = 0.3$, $p_2 = 0.23$, $r_1 = 0.15$, $r_2 = -0.1$, $a_1 = a_2 = 1$, $\alpha = 0.8$, $\beta = 0.5$).